

Partial summation of the nonlocal expansion for the gravitational effective action in 4 dimensions

A. G. Mirzabekian^a, G. A. Vilkovisky^{a,b}, V. V. Zhytnikov^{a,c}

^a*Lebedev Research Center in Physics, Leninsky Prospect 53, Moscow 117924, Russia*

^b*Lebedev Physics Institute, Academy of Sciences of Russia, Leninsky Prospect 53, Moscow 117924, Russia*

^c*Moscow State Pedagogical University, Davydovskii 4, Moscow 107140, Russia*

The vacuum action for the gravitational field admits a known expansion in powers of the Ricci tensor with nonlocal operator coefficients (form factors). We show that going over to a different basis of curvature invariants makes possible a partial summation of this expansion. Only the form factors of the Weyl-tensor invariants need be calculated. The full action is then uniquely recovered to all orders from the knowledge of the trace anomaly. We present an explicit expression for the partially summed action, and point out simplifications resulting in the vertex functions. An application to the effect of the vacuum gravitational waves is discussed.

Our starting point is the nonlocal expansion of the effective action for a given quantum field model in powers of a collective field strength [1-7]. In the purely gravitational sector this field strength boils down to the Ricci tensor R^μ_ν since the Riemann tensor gets expressed through R^μ_ν via the Bianchi identities [1-3]. For a generic field model the collective field strength is the set

$$\mathfrak{R} = (R^\mu_\nu, \hat{\mathcal{R}}_{\mu\nu}, \hat{P}) \quad (1)$$

which consists of the Ricci tensor, the commutator curvature $\hat{\mathcal{R}}_{\mu\nu} = [\nabla_\mu, \nabla_\nu]$ accounting for an arbitrary connection and satisfying the Jacobi identity, and the potential $\hat{P}$ (see refs. [2-7]). Both $\hat{\mathcal{R}}_{\mu\nu}$ and $\hat{P}$ are matrices in the space of fields disturbances. At every order in $\mathfrak{R}$ there is a finite basis of nonlocal curvature invariants [2], and the effective action is their linear combination. Denoting W the euclidean effective action with the term linear in the curvature subtracted, we have [1,2]

$$W = \sum_{n=2}^{\infty} \int dx g^{1/2} \sum_i \Gamma_i(\square_1, \dots \square_n, \square_{1+2}, \square_{2+3}, \dots) \mathfrak{R}_1 \dots \mathfrak{R}_n(i) \quad (2)$$

where, for a given order n , i labels a finite set of basis invariants, and $\mathfrak{R}_1 \dots \mathfrak{R}_n(i)$ in the n th-order basis invariant number i . For pure gravity, it has the structure

$$\mathfrak{R}_1 \dots \mathfrak{R}_n(i) = M_i(\nabla_1, \dots \nabla_n) R^\bullet_{\bullet 1} \dots R^\bullet_{\bullet n} \quad (3)$$

where $R^\bullet_{\bullet k} = R^\mu_\nu(x_k)$ is the Ricci tensor at a point x_k , $M_i(\nabla_1, \dots \nabla_n)$ is a monomial in covariant derivatives whose order does not exceed $2n$, ∇_k acts on $R^\bullet_{\bullet k}$, and, after the action of all operators, the points x_k are made coincident. The arguments of the form factors Γ_i in (2), $\square \equiv g^{\mu\nu} \nabla_\mu \nabla_\nu$, include all operators $\square_k$ acting on individual Ricci tensors $R^\bullet_{\bullet k}$ and all operators $\square_{k+p}$ acting on the products $R^\bullet_{\bullet k} R^\bullet_{\bullet p}$ of two Ricci tensors. For the full set of curvatures (1), the structure of basis invariants in the expansion (2) is similar.

The basis invariants are explicitly built in [2] to third order inclusive, and the respective form factors Γ_i are calculated in the one-loop approximation for a generic model of quantum fields [3-6]. The asymptotic behaviours of the form factors Γ_i at one of the arguments small and the others fixed [7] determine the effects of coherent [8,9] and noncoherent [10,11] vacuum radiation from the asymptotically flat systems.

When calculated by the dimensional regularization, the effective action of conformal invariant fields produces a local trace anomaly (for the original references and further developments see [5]). This property can also be formulated for a generic quantum field model with the inverse propagator of the form ¹

$$g^{\mu\nu}\nabla_\mu\nabla_\nu\hat{1} + (\hat{P} - \frac{1}{6}R\hat{1}) \quad (4)$$

provided that the effective action W is expressed in terms of the collective field strength (1). Assuming that, under the conformal transformation of all fields

$$\delta_\sigma = \int dx \sigma(x) \left(2g_{\mu\nu} \frac{\delta}{\delta g_{\mu\nu}} + \dots \right), \quad (5)$$

the commutator curvature and potential transform as follows

$$\delta_\sigma \hat{\mathcal{R}}_{\mu\nu} = 0, \quad \delta_\sigma \hat{P} = -2\sigma \hat{P}, \quad (6)$$

one obtains a local expression for $\delta_\sigma W$ in the universal form (eq. (28) below) [5,12].

Because the conformal transformation is inhomogeneous in the curvature, the expansion in powers of the curvature (2) does not preserve the exact conformal properties of the effective action. These properties can only be recovered order by order by an explicit calculation. To third order inclusive this calculation was done for the anomaly in two [4] and four [5] dimensions. The purpose of the present paper is turning this shortcoming into an advantage. We shall use the knowledge of the exact trace anomaly to perform a partial summation of the expansion (2).

As a first step we go over to a new basis of curvature invariants in which the gravitational field strength is represented, instead of R^μ_ν , by the set $C^\alpha_{\beta\gamma\sigma}, R$ consisting of the Weyl tensor and the Ricci scalar. To show that this set is complete, we use the corollary of the Bianchi identities for the Weyl tensor

$$\nabla^\gamma \nabla^\alpha C_{\alpha\beta\gamma\delta} = \frac{1}{2} \square R_{\beta\delta} - \frac{1}{2} \nabla_\gamma \nabla_\delta R^\gamma_\beta + \frac{1}{12} \nabla_\beta \nabla_\delta R - \frac{1}{12} g_{\beta\delta} \square R \quad (7)$$

¹Here and below, $\hat{1}$ is the unit matrix in the space of fields disturbances, and the sign conventions for the curvature are $R^\mu_{\alpha\nu\beta} = \partial_\nu \Gamma^\mu_{\alpha\beta} - \dots$, $R_{\alpha\beta} = R^\mu_{\alpha\mu\beta}$, $R = g^{\alpha\beta} R_{\alpha\beta}$.

to express R_ν^μ iteratively through $C_{\beta\gamma\sigma}^\alpha$ and R :

$$R_{\mu\nu} = C_{\mu\nu} + \frac{1}{3}\nabla_\mu\nabla_\nu\frac{1}{\square}R + \frac{1}{6}g_{\mu\nu}R + \mathcal{O}[\mathfrak{R}^2]. \quad (8)$$

Here

$$C_{\mu\nu} \stackrel{\text{def}}{=} \frac{2}{\square}\nabla^\beta\nabla^\alpha C_{\alpha\mu\beta\nu}, \quad (9)$$

and the relation inverse to (9) is ²

$$\begin{aligned} C_{\alpha\beta\gamma\delta} &= \frac{1}{\square}(\nabla_\alpha\nabla_\gamma C_{\beta\delta} + \nabla_\beta\nabla_\delta C_{\alpha\gamma} - \nabla_\beta\nabla_\gamma C_{\alpha\delta} - \nabla_\alpha\nabla_\delta C_{\beta\gamma}) \\ &\quad - \frac{1}{2}(g_{\alpha\gamma}C_{\beta\delta} + g_{\beta\delta}C_{\alpha\gamma} - g_{\beta\gamma}C_{\alpha\delta} - g_{\alpha\delta}C_{\beta\gamma}) + \mathcal{O}[\mathfrak{R}^2]. \end{aligned} \quad (10)$$

Expression (8) can now be used to rewrite the basis invariants (3) in terms of $C_{\beta\gamma\sigma}^\alpha$ and R . The result is that the action (2) becomes a sum of three contributions

$$W = W_C + W_{CR} + W_R \quad (11)$$

each of which is an expansion of the form (2), (3), and W_C contains the basis invariants with the Weyl tensor only:

$$\begin{aligned} W_C &= \sum_{n=2}^{\infty} \int dx g^{1/2} \sum_i \Gamma_i(\square_1, \dots, \square_n, \square_{1+2}, \square_{2+3}, \dots) \\ &\quad \times M_i(\nabla_1, \dots, \nabla_n) \mathbf{C}_1 \dots \mathbf{C}_n. \end{aligned} \quad (12)$$

Here $\mathbf{C}$ is the notation for $C_{\beta\gamma\sigma}^\alpha$ with the indices omitted, and we do not introduce a new notation for the form factors in (12) which are linear combinations of the form factors in (2). The two other terms in (11) are of a similar structure and contain the basis invariants either with both $C_{\beta\gamma\sigma}^\alpha$ and R (the term W_{CR}) or with R only (the term W_R).

The main assertion of the present paper is that the effective action W is completely determined by its term W_C and its trace anomaly. When the full set of field strengths (1) is present, the term W_C should include all invariants with $\mathbf{C}$, $\hat{\mathcal{R}}_{\mu\nu}$ and $\hat{P}$ but not R :

$$W_C = W_C(g_{\mu\nu}, \mathbf{C}, \hat{\mathcal{R}}_{\mu\nu}, \hat{P}). \quad (13)$$

²In lorentzian spacetime, the validity of relation (10) with $1/\square$ replaced by the retarded Green function $1/\square_{\text{ret}}$ [3] is limited to the in-vacuum state for gravitons (see also below).

The complementary contributions to W contain at least one Ricci scalar R in every term of expansion. Below we present (i) an explicit expression which completely recovers W given W_C , and (ii) a proof that this recovery is unique. We emphasize that the term W_C *is not conformal invariant*.³

For accomplishing the points (i) and (ii) above, we shall use the method of building conformal invariants proposed in [12,13]. The main tool is the solution

$$\phi(x|g) = 1 + \frac{1}{6} \left(\square - \frac{1}{6} R \right)^{-1} R \quad (14)$$

of the equation

$$\left(\square - \frac{1}{6} R \right) \phi = 0 \quad (15)$$

for a scalar field in an asymptotically flat euclidean spacetime with the boundary condition $\phi = 1$ at infinity. Under a conformal transformation that becomes the identity at infinity the solution transforms as follows

$$\phi(x|g_{\mu\nu} e^{2\sigma}) = e^{-\sigma(x)} \phi(x|g_{\mu\nu}), \quad (16)$$

and, therefore, the quantity

$$\bar{g}_{\mu\nu}(x) = \phi^2(x|g) g_{\mu\nu}(x) \quad (17)$$

is conformal invariant. Hence, any functional of $\bar{g}_{\mu\nu}(x)$ is conformal invariant. When used as a metric, $\bar{g}_{\mu\nu}(x)$ possesses the property that its Ricci scalar vanishes identically:

$$R(\bar{g}) \equiv 0. \quad (18)$$

On the other hand, ϕ in (14) admits an expansion of the same nature as the one used for the construction of the effective action (2):

$$\phi = 1 + \frac{1}{6} \frac{1}{\square} R + \frac{1}{36} \frac{1}{\square} \left(R \frac{1}{\square} R \right) + \dots \quad (19)$$

It is only in the framework of this expansion that the solution (14) will be used in the present context.

³The concept of conformal invariant part of the effective action is ill-defined. There are many different ways to decompose a given action into conformal invariant and anomalous parts, and there are many different actions producing one and the same trace anomaly (see [14] and references therein).

We are now able to prove that if two actions of the metric, $W'[g]$ and $W''[g]$, each in the form of expansion (2), or (11), give rise to one and the same trace

$$g_{\mu\nu} \frac{\delta W'}{\delta g_{\mu\nu}} = g_{\mu\nu} \frac{\delta W''}{\delta g_{\mu\nu}} \quad (20)$$

and have one and the same form factors of the Weyl-tensor invariants

$$W'_C[g] = W''_C[g], \quad (21)$$

then these actions coincide. Indeed, the difference $\Delta = W' - W''$ has the form of expansions (2) and (11) but with no pure-Weyl terms

$$\Delta[g] = \Delta_{CR}[g] + \Delta_R[g] \quad (22)$$

and is conformal invariant. Therefore, with $\bar{g}$ defined in (17), we have

$$\Delta[g] = \Delta[\bar{g}] = \Delta_{CR}[\bar{g}] + \Delta_R[\bar{g}] = 0 \quad (23)$$

where the vanishing takes place owing to eq. (18) and the fact that both Δ_{CR} and Δ_R contain at least one R in every term of expansion.

There remains to be presented an explicit expression for the full action W given W_C . This expression is of the following form:

$$\begin{aligned} W = & W_C(\phi^2 g_{\mu\nu}, \mathbf{C}, \hat{\mathcal{R}}_{\mu\nu}, \phi^{-2} \hat{P}) \\ & + \frac{1}{2(4\pi)^2} \int dx g^{1/2} \text{tr} \left\{ \frac{1}{1080} R^2 \hat{1} + \frac{1}{18} R \hat{P} \right. \\ & + \frac{1}{180} Z \left(C_{\alpha\beta\gamma\delta} C^{\alpha\beta\gamma\delta} + R_{\alpha\beta} R^{\alpha\beta} - \frac{1}{3} R^2 \right) \hat{1} \\ & + \frac{1}{180} R^{\alpha\beta} \nabla_\alpha Z \nabla_\beta Z \hat{1} + \frac{1}{360} (\nabla Z)^2 \left(-R + \square Z + \frac{1}{4} (\nabla Z)^2 \right) \hat{1} \\ & \left. + \frac{1}{12} Z \hat{\mathcal{R}}_{\mu\nu} \hat{\mathcal{R}}^{\mu\nu} + \frac{1}{2} Z \hat{P} \hat{P} \right\} \end{aligned} \quad (24)$$

where tr denotes the matrix trace, Z is the following quantity

$$Z = \ln \phi^2, \quad (25)$$

and ϕ is defined in (14). The first term in (24) is built out of the original W_C in (13) by replacing the metric $g_{\mu\nu}$ *wherever it appears* with $\phi^2 g_{\mu\nu}$, and $\hat{P}$

with $\phi^{-2}\hat{P}$. Specifically, since in the purely gravitational sector W_C is given by expansion (12), the respective contribution to (24) is of the form

$$W_C(\phi^2 g_{\mu\nu}, \mathbf{C}) = \sum_{n=2}^{\infty} \int dx \bar{g}^{1/2} \sum_i \Gamma_i(\bar{\square}_1, \dots, \bar{\square}_n, \bar{\square}_{1+2}, \bar{\square}_{2+3}, \dots) \\ \times M_i(\bar{\nabla}_1, \dots, \bar{\nabla}_n) \mathbf{C}_1 \dots \mathbf{C}_n \quad (26)$$

where all symbols with the bar refer to the metric $\bar{g}_{\mu\nu}$ in (17). The explicit addition to W_C in (24) contains Z up to the fourth power but, if needed, the power can be reduced by using the equation

$$\square Z + \frac{1}{2}(\nabla Z)^2 = \frac{1}{3}R \quad (27)$$

that Z in (25) satisfies.

To prove that expression (24) for W correctly recovers the original action, one must first show that all pure-Weyl invariants of W are contained in the original W_C . This is easily seen upon the insertion of expansion (19) in (25) and (24). Next, one must show that the action (24) gives rise to the correct trace anomaly. Since the term with W_C in (24) is conformal invariant by construction, the anomaly is produced by the remaining explicit terms. The direct calculation using (6) with δ_σ in (5) gives

$$-\delta_\sigma W = \frac{1}{(4\pi)^2} \int dx g^{1/2} \sigma(x) \text{tr } \hat{a}_2(x, x), \quad (28)$$

$$\hat{a}_2(x, x) = \frac{1}{180} \square R \hat{1} + \frac{1}{6} \square \hat{P} + \frac{1}{180} \left(C_{\alpha\beta\gamma\delta} C^{\alpha\beta\gamma\delta} + R_{\alpha\beta} R^{\alpha\beta} - \frac{1}{3} R^2 \right) \hat{1} \\ + \frac{1}{12} \hat{\mathcal{R}}_{\mu\nu} \hat{\mathcal{R}}^{\mu\nu} + \frac{1}{2} \hat{P} \hat{P}. \quad (29)$$

The latter quantity is the second DeWitt coefficient [15] at coincident points for the operator (4). This is the correct expression for the trace anomaly in four dimensions (see [5,12] and references therein).

The significance of the result above is in the fact that it suffices to calculate only the form factors of the basis invariants with the Weyl tensor (and matter field strengths); they accumulate the dynamical information and are generally not controlled by symmetries. The remaining form factors in the

effective action are then obtained automatically by inserting expansion (19) in (24). Specifically, if the form factors of the Weyl tensor are known to third order inclusive (as is presently the case with the one-loop effective action [3-7]), then all contributions to W containing two or three Weyl tensors and any number of Ricci scalars are also known. The contributions of invariants with the Ricci scalar only are all known in advance in the closed form. This is a partial summation of the series (2) for the effective action.

The transition to the $C_{\beta\gamma\sigma}^\alpha$, R basis simplifies greatly the form factors obtained in [5,6]. The one-loop effective action for the operator (4) takes the form

$$\begin{aligned}
-W = & \frac{1}{2(4\pi)^2} \int dx g^{1/2} \text{tr} \left\{ -\frac{1}{1080} R^2 \hat{1} - \frac{1}{18} R \hat{P} \right. \\
& - \frac{1}{120} C_{\alpha\beta\gamma\delta} \left(\ln(-\square/\mu^2) - \frac{16}{15} \right) C^{\alpha\beta\gamma\delta} \hat{1} \\
& - \frac{1}{12} \hat{\mathcal{R}}_{\mu\nu} \left(\ln(-\square/\mu^2) - \frac{2}{3} \right) \hat{\mathcal{R}}^{\mu\nu} - \frac{1}{2} \hat{P} \left(\ln(-\square/\mu^2) \right) \hat{P} \\
& \left. + \sum_i \Gamma_i(-\square_1, -\square_2, -\square_3) \overline{\mathfrak{R}_1 \mathfrak{R}_2 \mathfrak{R}_3}(i) + \mathcal{O}[\mathfrak{R}^4] \right\} \quad (30)
\end{aligned}$$

where the only arbitrary parameter is $\mu^2 > 0$, and the new cubic invariants $\overline{\mathfrak{R}_1 \mathfrak{R}_2 \mathfrak{R}_3}(i)$ are obtained from the old ones $\mathfrak{R}_1 \mathfrak{R}_2 \mathfrak{R}_3(i)$ in [5,6] by inserting the expression (8) for $R_{\mu\nu}$. As a result, the full list of new cubic invariants differs from the old one only by the replacement of $R_{\mu\nu}$ with $C_{\mu\nu}$ (eq. (9)) but the form factors of the invariants with the Ricci scalar change drastically. In the purely gravitational sector there are five such invariants

$$\begin{aligned}
\overline{\mathfrak{R}_1 \mathfrak{R}_2 \mathfrak{R}_3}(9) &= R_1 R_2 R_3 \hat{1}, \quad \overline{\mathfrak{R}_1 \mathfrak{R}_2 \mathfrak{R}_3}(11) = C_1^{\mu\nu} C_{2\mu\nu} R_3 \hat{1}, \\
\overline{\mathfrak{R}_1 \mathfrak{R}_2 \mathfrak{R}_3}(22) &= C_1^{\alpha\beta} \nabla_\alpha R_2 \nabla_\beta R_3 \hat{1}, \quad \overline{\mathfrak{R}_1 \mathfrak{R}_2 \mathfrak{R}_3}(23) = \nabla^\mu C_1^{\nu\alpha} \nabla_\nu C_{2\mu\alpha} R_3 \hat{1}, \\
\overline{\mathfrak{R}_1 \mathfrak{R}_2 \mathfrak{R}_3}(27) &= \nabla_\alpha \nabla_\beta C_1^{\mu\nu} \nabla_\mu \nabla_\nu C_2^{\alpha\beta} R_3 \hat{1}
\end{aligned}$$

whose form factors were the most complicated ones in the table of ref. [6]. Now they factorize into elementary functions of two variables and take the form

$$\Gamma_9 = \frac{1}{19440} \left(\frac{2}{\square_3} - \frac{\square_1}{\square_2 \square_3} \right), \quad (31)$$

$$\begin{aligned}\Gamma_{11} = & \frac{1}{540} \left(\frac{4}{\square_2} - \frac{1}{\square_3} - \frac{2\square_1}{\square_2\square_3} - \frac{\square_3}{\square_1\square_2} \right) + \frac{1}{720} \left(\frac{10\square_1}{\square_2} \right. \\ & \left. + \frac{6\square_1}{\square_3} - \frac{2\square_1^2}{\square_2\square_3} - \frac{14\square_3}{\square_2} + \frac{3\square_3^2}{\square_1\square_2} \right) \frac{\ln(\square_1/\square_2)}{(\square_1 - \square_2)},\end{aligned}\quad (32)$$

$$\Gamma_{22} = \frac{1}{1620\square_2\square_3}, \quad (33)$$

$$\begin{aligned}\Gamma_{23} = & \frac{1}{135} \left(\frac{1}{\square_1\square_2} - \frac{2}{\square_2\square_3} \right) \\ & + \frac{1}{180} \left(\frac{8}{\square_2} - \frac{2\square_1}{\square_2\square_3} - \frac{3\square_3}{\square_1\square_2} \right) \frac{\ln(\square_1/\square_2)}{(\square_1 - \square_2)},\end{aligned}\quad (34)$$

$$\Gamma_{27} = \frac{-1}{135\square_1\square_2\square_3} + \frac{1}{180} \left(\frac{3}{\square_1\square_2} - \frac{2}{\square_2\square_3} \right) \frac{\ln(\square_1/\square_2)}{(\square_1 - \square_2)}. \quad (35)$$

Furthermore, upon the replacement $g_{\mu\nu} \rightarrow \bar{g}_{\mu\nu} = \phi^2 g_{\mu\nu}$ in the quadratic terms of (30), the contributions proportional to

$$\frac{\ln(\square_1/\square_2)}{(\square_1 - \square_2)} \quad (36)$$

in (31)–(35) get *exactly absorbed* by the second-order form factors

$$\ln(-\square/\mu^2) \rightarrow \ln(-\bar{\square}/\mu^2). \quad (37)$$

The remaining tree contributions in (31)–(35) give rise to the trace anomaly with accuracy $\mathcal{O}[\Re^3]$ [5], and, as one can check, they are the expansion terms of the exact action in (24). Thus expression (24) is in complete agreement with the results of explicit calculations.

The advantages provided by the transition to the $C_{\beta\gamma\sigma}^\alpha, R$ basis and by the partial summation accomplished in (24) have an immediate impact on applications. The complicated check [9] that the $1/\square, \square \rightarrow 0$, asymptotic terms of the form factors cancel in the energy-momentum tensor is no more needed since this cancellation becomes manifest. The contribution to the energy of the vacuum gravitational waves that was caused by the $1/\square$ asymptotic behaviours [8,9] is now generated by a different, much simpler mechanism.

Following the lines of ref. [9], one must now introduce a new generalized current, $I^{\mu\alpha\nu\beta}(-\square, x)$, conjugate to the variation of the Weyl tensor

$$\delta W = \frac{1}{2(4\pi)^2} \int dx g^{1/2} I^{\mu\alpha\nu\beta}(-\square, x) \delta C_{\mu\alpha\nu\beta}, \quad (38)$$

and it turns out that the $O(\square^0)$ term of $I^{\mu\alpha\nu\beta}(-\square, x)$ gives directly the quantity $K^{\mu\alpha\nu\beta}(x)$ which in refs. [8,9] determined the Bondi-Sachs news functions $\mathcal{C}_1, \mathcal{C}_2$ of the vacuum gravitational waves

$$\frac{\partial}{\partial u}(\mathcal{C}_1 + i\mathcal{C}_2) = \frac{\partial}{\partial u} \left\{ \frac{r}{8\pi} m_\mu m_\nu \frac{1}{\square_{\text{ret}}} \nabla_\alpha \nabla_\beta K^{[\mu\alpha][\nu\beta]}(x) \right\} \Big|_{\mathcal{I}^+}. \quad (39)$$

Here u is the retarded time, m is the complex vector of the null tetrad, and $r \rightarrow \infty$ is the luminosity distance to the future null infinity $\mathcal{I}^+$.

To present an example, we pick out the sector of the effective action linear in the potential $\hat{P}$. The $1/\square, \square \rightarrow 0$, behaviours of the form factors in this sector, present in the expansion over the R_ν^μ basis [7,9], are now absent completely. On the other hand, for the contribution linear in $\hat{P}$ to the vacuum news functions (39) we obtain straight away

$$K^{\mu\alpha\nu\beta} = \frac{1}{3} \text{tr} \frac{1}{(\square_2 - \square_3)^2} \left(3\square_2 - \square_3 - 2\square_2^2 \frac{\ln(\square_2/\square_3)}{(\square_2 - \square_3)} \right) C_2^{\mu\alpha\nu\beta} \hat{P}_3. \quad (40)$$

The latter expression agrees with the result in [9] and is of a simpler form.

Finally, one more advantage of the $C_{\beta\gamma\sigma}^\alpha, R$ basis arises from the fact that, in lorentzian spacetime, the use of relation (8) imposes no limitation on the initial state. There is only one solution of eq. (7) for $R_{\beta\delta}$ with the appropriate decrease at $\mathcal{I}^\pm$, and this solution is generated by the retarded Green function for any choice of the in-state. In contrast with this case, the use of the retarded Green function in the solution of the Bianchi identities for the Riemann tensor [1-3] excludes an incoming gravitational wave. This concerns also eq. (10) above. Normally, the effective action calculated for a given vacuum remains valid for all coherent states based on this vacuum. However, because in the calculations of refs. [3-7] the Riemann tensor was excluded via the Bianchi identities, the results of these calculations are valid only for the vacuum state. The effective action for coherent graviton in-states cannot be expanded over the R_ν^μ basis at all but the $C_{\beta\gamma\sigma}^\alpha, R$ basis is suitable for this purpose since the only relation involved is eq. (8).

The present work was supported in part by the Russian Foundation for Fundamental Research Grant 93-02-15594, INTAS Grant 93-0493, and the joint Grant MQY300 of the International Science Foundation and Russian Government.

References

- [1] G. A. Vilkovisky, *Class. Quantum Grav.* 9 (1992) 895; preprint CERN-TH.6392/92; *Publ. Inst. Rech. Math. Avancée, R.C.P. 25, Vol. 43* (Strasbourg, 1992) p. 203.
- [2] A. O. Barvinsky, Yu. V. Gusev, G. A. Vilkovisky and V. V. Zhytnikov, *J. Math. Phys.* 35 (1994) 3525.
- [3] A. O. Barvinsky and G. A. Vilkovisky, *Nucl. Phys.* B282 (1987) 163; B333 (1990) 471, 512.
- [4] A. O. Barvinsky, Yu. V. Gusev, G. A. Vilkovisky and V. V. Zhytnikov, *J. Math. Phys.* 35 (1994) 3543.
- [5] A. O. Barvinsky, Yu. V. Gusev, G. A. Vilkovisky and V. V. Zhytnikov, *Nucl. Phys.* B439 (1995) 561.
- [6] A. O. Barvinsky, Yu. V. Gusev, V. V. Zhytnikov and G. A. Vilkovisky, *Covariant perturbation theory (IV). Third order in the curvature*, University of Manitoba at Winnipeg Report (1993) pp. 1–192.
- [7] A. O. Barvinsky, Yu. V. Gusev, V. V. Zhytnikov and G. A. Vilkovisky, *Class. Quantum Grav.* 12 (1995) 2157.
- [8] A. G. Mirzabekian and G. A. Vilkovisky, *Gravitational waves generated by the vacuum stress*, *Phys. Rev. Lett.* (1995) to appear.
- [9] A. G. Mirzabekian and G. A. Vilkovisky, *Class. Quantum Grav.* 12 (1995) 2173.
- [10] A. G. Mirzabekian and G. A. Vilkovisky, *Phys. Lett. B* 317 (1993) 517.
- [11] A. G. Mirzabekian, *Zh. Eksp. Teor. Fiz.* 106 (1994) 5 [*JETP* 79 (1994) 1].

- [12] E. S. Fradkin and G. A. Vilkovisky, Phys. Lett. B 73 (1978) 209.
- [13] G. A. Vilkovisky, *in* Quantum theory of gravity, ed. S. M. Christensen (Hilger, Bristol, 1984).
- [14] A. O. Barvinsky, A. G. Mirzabekian and V. V. Zhytnikov, Proc. 6th Quantum Gravity Seminar, Moscow 1995, to be published.
- [15] A. O. Barvinsky and G. A. Vilkovisky, Phys. Rep. 119 (1985) 1.